




Bunions are a **progressive disorder** and do not go away on their own without surgical correction.<sup>1</sup>

About  
**1 in 4**  
US adults  
suffer from  
bunions.<sup>2</sup>



**Up to 30%** of patients  
with bunions also have MTA.<sup>3,4</sup>



**Adductoplasty**<sup>®</sup>  
Midfoot Correction System

**Lapiplasty**<sup>®</sup>  
3D Bunion Correction

Individual results may vary depending on age, weight, health and other variables.

There are risks and recovery takes time.

Risks include infection, pain, implant loosening and loss of correction with improper bone healing. If any of these occur, additional treatments may be needed.

For more information on benefits, risks and recovery, visit [Lapiplasty.com](http://Lapiplasty.com).

1. American College of Foot and Ankle Surgeons (ACFAS) Website ©2022
2. Nix S, et al. J Foot Ankle Res. 2010. 27:3:21.
3. Aiyer AA, et al. FAI 2016; 37: 165-171.
4. Aiyer AA, et al. FAI. 2014; 35: 1292-1297.

**TREACE**<sup>®</sup>  
Medical Concepts, Inc.  
*The Leader in Hallux Valgus Surgery™*

Registered marks and trademarks of Treace Medical Concepts, Inc.  
Pat. [treace.com/patents](http://treace.com/patents)  
©2024 Treace Medical Concepts, Inc. All Rights Reserved. M2104B

**Adductoplasty**<sup>®</sup>  
Midfoot Correction System

**Because not all  
bunions are the same.**




**Complementing Lapiplasty<sup>®</sup> 3D Bunion Correction.**

# Know Your Bunions and Your Options

## What is Metatarsus Adductus (MTA)?


MTA is a common foot deformity that causes the front half of the foot, or forefoot, to turn inward. A full correction of the bunion may require repositioning the bones of the 2nd and 3rd toes in order to make room for complete first toe realignment. Failure to address MTA increases the risk of bunion recurrence.<sup>3</sup>



Midfoot Malalignment  
Blocks Bunion Correction

## Why have Adductoplasty®?

The **Adductoplasty® Procedure** corrects the alignment of the 2nd and 3rd toes, creating the necessary space to perform the **Lapiplasty® Procedure** and bring the big toe back into normal alignment.



Midfoot Alignment  
Allows Full Bunion Correction

## The Lapiplasty® Family of Procedures

Not all bunions are the same. Sometimes bunions are accompanied by big toe arthritis or a midfoot deformity like MTA. Adductoplasty® instrumentation, specifically designed to correct this deformity, along with the associated implants, provide a comprehensive system that allows for realignment and stabilization of the midfoot.



Bunion without MTA



Correction with Lapiplasty®  
3D Bunion Correction



Bunion with MTA



Correction with  
Adductoplasty® and  
Lapiplasty® Procedure